

TABLETS

DOLOBID®
(DIFLUNISAL)

Cardiovascular Risk


- NSAIDS may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk. (See **WARNINGS**.)
- DOLOBID is contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery (see **WARNINGS**).

Gastrointestinal Risk

- NSAIDS cause an increased risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious gastrointestinal events. (See **WARNINGS**.)

DESCRIPTION

Diflunisal is 2', 4'-difluoro-4-hydroxy-3-biphenylcarboxylic acid. Its empirical formula is C₁₃H₈F₂O₃ and its structural formula is:



Diflunisal has a molecular weight of 250.20. It is a stable, white, crystalline compound with a melting point of 211-213°C. It is practically insoluble in water at neutral or acidic pH. Because it is an organic acid, it dissolves readily in dilute alkali to give a moderately stable solution at room temperature. It is soluble in most organic solvents including ethanol, methanol, and acetone.

DOLOBID® (Diflunisal) is available in 250 and 500 mg tablets for oral administration. Tablets DOLOBID contain the following inactive ingredients: cellulose, FD&C Yellow 6, hydroxypropyl cellulose, hydroxypropyl methylcellulose, magnesium stearate, starch, talc, and titanium dioxide.

CLINICAL PHARMACOLOGY

Action

DOLOBID is a non-steroidal drug with analgesic, anti-inflammatory and antipyretic properties. It is a peripherally-acting non-narcotic analgesic drug. Habituation, tolerance and addiction have not been reported.

Diflunisal is a difluorophenyl derivative of salicylic acid. Chemically, diflunisal differs from aspirin (acetylsalicylic acid) in two respects. The first of these two is the presence of a difluorophenyl substituent at carbon 1. The second difference is the removal of the O-acetyl group from the carbon 4 position. Diflunisal is not metabolized to salicylic acid, and the fluorine atoms are not displaced from the difluorophenyl ring structure.

The precise mechanism of the analgesic and anti-inflammatory actions of diflunisal is not known. Diflunisal is a prostaglandin synthetase inhibitor. In animals, prostaglandins sensitize afferent nerves and

potentiate the action of bradykinin in inducing pain. Since prostaglandins are known to be among the mediators of pain and inflammation, the mode of action of diflunisal may be due to a decrease of prostaglandins in peripheral tissues.

Pharmacokinetics and Metabolism

DOLOBID is rapidly and completely absorbed following oral administration with peak plasma concentrations occurring between 2 to 3 hours. The drug is excreted in the urine as two soluble glucuronide conjugates accounting for about 90% of the administered dose. Little or no diflunisal is excreted in the feces. Diflunisal appears in human milk in concentrations of 2-7% of those in plasma. More than 99% of diflunisal in plasma is bound to proteins.

As is the case with salicylic acid, concentration-dependent pharmacokinetics prevail when DOLOBID is administered; a doubling of dosage produces a greater than doubling of drug accumulation. The effect becomes more apparent with repetitive doses. Following single doses, peak plasma concentrations of 41 ± 11 $\mu\text{g/mL}$ (mean \pm S.D.) were observed following 250 mg doses, 87 ± 17 $\mu\text{g/mL}$ were observed following 500 mg and 124 ± 11 $\mu\text{g/mL}$ following single 1000 mg doses. However, following administration of 250 mg b.i.d., a mean peak level of 56 ± 14 $\mu\text{g/mL}$ was observed on day 8, while the mean peak level after 500 mg b.i.d. for 11 days was 190 ± 33 $\mu\text{g/mL}$. In contrast to salicylic acid which has a plasma half-life of 2½ hours, the plasma half-life of diflunisal is 3 to 4 times longer (8 to 12 hours), because of a difluorophenyl substituent at carbon 1. Because of its long half-life and nonlinear pharmacokinetics, several days are required for diflunisal plasma levels to reach steady state following multiple doses. For this reason, an initial loading dose is necessary to shorten the time to reach steady state levels, and 2 to 3 days of observation are necessary for evaluating changes in treatment regimens if a loading dose is not used.

Studies in baboons to determine passage across the blood-brain barrier have shown that only small quantities of diflunisal, under normal or acidotic conditions are transported into the cerebrospinal fluid (CSF). The ratio of blood/CSF concentrations after intravenous doses of 50 mg/kg or oral doses of 100 mg/kg of diflunisal was 100:1. In contrast, oral doses of 500 mg/kg of aspirin resulted in a blood/CSF ratio of 5:1.

Mild to Moderate Pain

DOLOBID is a peripherally-acting analgesic agent with a long duration of action. DOLOBID produces significant analgesia within 1 hour and maximum analgesia within 2 to 3 hours.

Consistent with its long half-life, clinical effects of DOLOBID mirror its pharmacokinetic behavior, which is the basis for recommending a loading dose when instituting therapy. Patients treated with DOLOBID, on the first dose, tend to have a slower onset of pain relief when compared with drugs achieving comparable peak effects. However, DOLOBID produces longer-lasting responses than the comparative agents.

Comparative single dose clinical studies have established the analgesic efficacy of DOLOBID at various dose levels relative to other analgesics. Analgesic effect measurements were derived from hourly evaluations by patients during eight and twelve-hour postdosing observation periods. The following information may serve as a guide for prescribing DOLOBID.

DOLOBID 500 mg was comparable in analgesic efficacy to aspirin 650 mg, acetaminophen 600 mg or 650 mg, and acetaminophen 650 mg with propoxyphene napsylate 100 mg. Patients treated with DOLOBID had longer lasting responses than the patients treated with the comparative analgesics.

DOLOBID 1000 mg was comparable in analgesic efficacy to acetaminophen 600 mg with codeine 60 mg. Patients treated with DOLOBID had longer lasting responses than the patients who received acetaminophen with codeine.

A loading dose of 1000 mg provides faster onset of pain relief, shorter time to peak analgesic effect, and greater peak analgesic effect than an initial 500 mg dose.

In contrast to the comparative analgesics, a significantly greater proportion of patients treated with DOLOBID did not re-medicate and continued to have a good analgesic effect eight to twelve hours after dosing. Seventy-five percent (75%) of patients treated with DOLOBID continued to have a good analgesic response at four hours. When patients having a good analgesic response at four hours were followed, 78% of these patients continued to have a good analgesic response at eight hours and 64% at twelve hours.

Chronic Anti-inflammatory Therapy in Osteoarthritis and Rheumatoid Arthritis

In the controlled, double-blind clinical trials in which DOLOBID (500 mg to 1000 mg a day) was compared with anti-inflammatory doses of aspirin (2-4 grams a day), patients treated with DOLOBID had a significantly lower incidence of tinnitus and of adverse effects involving the gastrointestinal system than patients treated with aspirin. (See also *Effect on Fecal Blood Loss*).

Osteoarthritis

The effectiveness of DOLOBID for the treatment of osteoarthritis was studied in patients with osteoarthritis of the hip and/or knee. The activity of DOLOBID was demonstrated by clinical improvement in the signs and symptoms of disease activity.

In a double-blind multicenter study of 12 weeks' duration in which dosages were adjusted according to patient response, DOLOBID, 500 or 750 mg daily, was shown to be comparable in effectiveness to aspirin, 2000 or 3000 mg daily. In open-label extensions of this study to 24 or 48 weeks, DOLOBID continued to show similar effectiveness and generally was well tolerated.

Rheumatoid Arthritis

In controlled clinical trials, the effectiveness of DOLOBID was established for both acute exacerbations and long-term management of rheumatoid arthritis. The activity of DOLOBID was demonstrated by clinical improvement in the signs and symptoms of disease activity.

In a double-blind multicenter study of 12 weeks' duration in which dosages were adjusted according to patient response, DOLOBID 500 or 750 mg daily was comparable in effectiveness to aspirin 2600 or 3900 mg daily. In open-label extensions of this study to 52 weeks, DOLOBID continued to be effective and was generally well tolerated.

DOLOBID 500, 750, or 1000 mg daily was compared with aspirin 2000, 3000, or 4000 mg daily in a multicenter study of 8 weeks' duration in which dosages were adjusted according to patient response. In this study, DOLOBID was comparable in efficacy to aspirin.

In a double-blind multicenter study of 12 weeks' duration in which dosages were adjusted according to patient needs, DOLOBID 500 or 750 mg daily and ibuprofen 1600 or 2400 mg daily were comparable in effectiveness and tolerability.

In a double-blind multicenter study of 12 weeks' duration, DOLOBID 750 mg daily was comparable in efficacy to naproxen 750 mg daily. The incidence of gastrointestinal adverse effects and tinnitus was comparable for both drugs. This study was extended to 48 weeks on an open-label basis. DOLOBID continued to be effective and generally well tolerated.

In patients with rheumatoid arthritis, DOLOBID and gold salts may be used in combination at their usual dosage levels. In clinical studies, DOLOBID added to the regimen of gold salts usually resulted in additional symptomatic relief but did not alter the course of the underlying disease.

Antipyretic Activity

DOLOBID is not recommended for use as an antipyretic agent. In single 250 mg, 500 mg, or 750 mg doses, DOLOBID produced measurable but not clinically useful decreases in temperature in patients with fever; however, the possibility that it may mask fever in some patients, particularly with chronic or high doses, should be considered.

Uricosuric Effect

In normal volunteers, an increase in the renal clearance of uric acid and a decrease in serum uric acid was observed when DOLOBID was administered at 500 mg or 750 mg daily in divided doses. Patients on long-term therapy taking DOLOBID at 500 mg to 1000 mg daily in divided doses showed a prompt and consistent reduction across studies in mean serum uric acid levels, which were lowered as much as 1.4 mg%. It is not known whether DOLOBID interferes with the activity of other uricosuric agents.

Effect on Platelet Function

As an inhibitor of prostaglandin synthetase, DOLOBID has a dose-related effect on platelet function and bleeding time. In normal volunteers, 250 mg b.i.d. for 8 days had no effect on platelet function, and 500 mg b.i.d., the usual recommended dose, had a slight effect. At 1000 mg b.i.d., which exceeds the maximum recommended dosage, however, DOLOBID inhibited platelet function. In contrast to aspirin, these effects of DOLOBID were reversible, because of the absence of the chemically labile and biologically reactive 0-acetyl group at the carbon 4 position. Bleeding time was not altered by a dose of 250 mg b.i.d., and was only slightly increased at 500 mg b.i.d. At 1000 mg b.i.d., a greater increase occurred, but was not statistically significantly different from the change in the placebo group.

Effect on Fecal Blood Loss

When DOLOBID was given to normal volunteers at the usual recommended dose of 500 mg twice daily, fecal blood loss was not significantly different from placebo. Aspirin at 1000 mg four times daily produced the expected increase in fecal blood loss. DOLOBID at 1000 mg twice daily (NOTE: exceeds the recommended dosage) caused a statistically significant increase in fecal blood loss, but this increase was only one-half as large as that associated with aspirin 1300 mg twice daily.

Effect on Blood Glucose

DOLOBID did not affect fasting blood sugar in diabetic patients who were receiving tolbutamide or placebo.

INDICATIONS AND USAGE

Carefully consider the potential benefits and risks of DOLOBID and other treatment options before deciding to use DOLOBID. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

DOLOBID is indicated for acute or long-term use for symptomatic treatment of the following:

1. Mild to moderate pain
2. Osteoarthritis
3. Rheumatoid arthritis

CONTRAINDICATIONS

DOLOBID is contraindicated in patients with known hypersensitivity to diflunisal or the excipients (see **DESCRIPTION**).

DOLOBID should not be given to patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Severe, rarely fatal, anaphylactic/anaphylactoid reactions to NSAIDs have been reported in such patients (see **WARNINGS – Anaphylactic/Anaphylactoid Reactions**, and **PRECAUTIONS – Preexisting asthma**).

DOLOBID is contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery (see **WARNINGS**).

WARNINGS

CARDIOVASCULAR EFFECTS

Cardiovascular Thrombotic Events

Clinical trials of several COX-2 selective and nonselective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, myocardial infarction, and stroke, which can be fatal. All NSAIDs, both COX-2 selective and nonselective, may have a similar risk. Patients with known CV disease or risk factors for CV disease may be at greater risk. To minimize the potential risk for an adverse CV event in patients treated with an NSAID, the lowest effective dose should be used for the shortest duration possible. Physicians and patients should remain alert for the development of such events, even in the absence of previous CV symptoms. Patients should be informed about the signs and/or symptoms of serious CV events and the steps to take if they occur.

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID does increase the risk of serious GI events (see **GI WARNINGS**).

Two large, controlled, clinical trials of a COX-2 selective NSAID for the treatment of pain in the first 10-14 days following CABG surgery found an increased incidence of myocardial infarction and stroke (see **CONTRAINDICATIONS**).

Hypertension

NSAIDs, including DOLOBID, can lead to onset of new hypertension or worsening of pre-existing hypertension, either of which may contribute to the increased incidence of CV events. Patients taking thiazides or loop diuretics may have impaired response to these therapies when taking NSAIDs. NSAIDs, including DOLOBID, should be used with caution in patients with hypertension. Blood pressure (BP) should be monitored closely during the initiation of NSAID treatment and throughout the course of therapy.

Congestive Heart Failure and Edema

Fluid retention and edema have been observed in some patients taking NSAIDs. DOLOBID should be used with caution in patients with fluid retention or heart failure.

Gastrointestinal Effects – Risk of Ulceration, Bleeding, and Perforation

NSAIDs, including DOLOBID, can cause serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients, who develop a serious upper GI adverse event on NSAID therapy is symptomatic. Upper GI ulcers, gross bleeding, or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3-6 months, and in about 2-4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of

developing a serious GI event at some time during the course of therapy. However, even short-term therapy is not without risk.

NSAIDs should be prescribed with extreme caution in those with prior history of ulcer disease or gastrointestinal bleeding. Patients with a *prior history of peptic ulcer disease and/or gastrointestinal bleeding* who use NSAIDs have a greater than 10-fold increased risk for developing a GI bleed compared to patients with neither of these risk factors. Other factors that increase the risk for GI bleeding in patients treated with NSAIDs include concomitant use of oral corticosteroids or anticoagulants, longer duration of NSAID therapy, smoking, use of alcohol, older age, and poor general health status. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and therefore, special care should be taken in treating this population.

To minimize the potential risk for an adverse GI event in patients treated with an NSAID, the lowest effective dose should be used for the shortest possible duration. Patients and physicians should remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected. This should include discontinuation of the NSAID until a serious GI adverse event is ruled out. For high risk patients, alternate therapies that do not involve NSAIDs should be considered.

Renal Effects

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of a nonsteroidal anti-inflammatory drug may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors, patients who are volume-depleted, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state.

Advanced Renal Disease

No information is available from controlled clinical studies regarding the use of DOLOBID in patients with advanced renal disease. Therefore, treatment with DOLOBID is not recommended in these patients with advanced renal disease. If DOLOBID therapy must be initiated, close monitoring of the patient's renal function is advisable.

Anaphylactic/Anaphylactoid Reactions

As with other NSAIDs, anaphylactic/anaphylactoid reactions may occur in patients without known prior exposure to DOLOBID. DOLOBID should not be given to patients with the aspirin triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking aspirin or other NSAIDs (see **CONTRAINDICATIONS** and **PRECAUTIONS – Preexisting Asthma**). Emergency help should be sought in cases where an anaphylactic/anaphylactoid reaction occurs.

Skin Reactions

NSAIDs, including DOLOBID, can cause serious skin adverse events such as exfoliative dermatitis, Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. These serious events may occur without warning. Patients should be informed about the signs and symptoms of serious skin manifestations and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

Hypersensitivity Syndrome

A potentially life-threatening, apparent hypersensitivity syndrome has been reported. This multisystem syndrome includes constitutional symptoms (fever, chills), and cutaneous findings (see **ADVERSE REACTIONS, Dermatologic**). It may also include involvement of major organs (changes in liver function, jaundice, leukopenia, thrombocytopenia, eosinophilia, disseminated intravascular coagulation, renal impairment, including renal failure), and less specific findings (adenitis, arthralgia, myalgia, arthritis, malaise, anorexia, disorientation). If evidence of hypersensitivity occurs, therapy with DOLOBID should be discontinued.

Pregnancy

In late pregnancy, as with other NSAIDs, DOLOBID should be avoided because it may cause premature closure of the ductus arteriosus.

PRECAUTIONS

General

DOLOBID cannot be expected to substitute for corticosteroids or to treat corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to disease exacerbation. Patients on prolonged corticosteroid therapy should have their therapy tapered slowly if a decision is made to discontinue corticosteroids.

The pharmacological activity of DOLOBID in reducing fever and inflammation may diminish the utility of these diagnostic signs in detecting complications of presumed noninfectious, painful conditions.

Hepatic Effects

Borderline elevations of one or more liver tests may occur in up to 15% of patients taking NSAIDs including DOLOBID. These laboratory abnormalities may progress, may remain unchanged, or may be transient with continuing therapy. Notable elevations of ALT or AST (approximately three or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials with NSAIDs. In addition, rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis and hepatic failure, some of them with fatal outcomes have been reported.

A patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of a more severe hepatic reaction while on therapy with DOLOBID. If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash, etc.), DOLOBID should be discontinued.

Hematological Effects

Anemia is sometimes seen in patients receiving NSAIDs, including DOLOBID. This may be due to fluid retention, occult or gross GI blood loss, or an incompletely described effect upon erythropoiesis. Patients on long-term treatment with NSAIDs, including DOLOBID, should have their hemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia.

NSAIDs inhibit platelet aggregation and have been shown to prolong bleeding time in some patients. Unlike aspirin, their effect on platelet function is quantitatively less, of shorter duration, and reversible. Patients receiving DOLOBID who may be adversely affected by alterations in platelet function, such as those with coagulation disorders or patients receiving anticoagulants, should be carefully monitored.

Preexisting Asthma

Patients with asthma may have aspirin-sensitive asthma. The use of aspirin in patients with aspirin-sensitive asthma has been associated with severe bronchospasm which can be fatal. Since cross reactivity, including bronchospasm, between aspirin and other nonsteroidal anti-inflammatory drugs has been reported in such aspirin-sensitive patients, DOLOBID should not be administered to patients with this form of aspirin sensitivity and should be used with caution in patients with preexisting asthma.

Ocular Effects

Because of reports of adverse eye findings with agents of this class, it is recommended that patients who develop eye complaints during treatment with DOLOBID have ophthalmologic studies.

Reye's Syndrome

Acetylsalicylic acid has been associated with Reye's syndrome. Because diflunisal is a derivative of salicylic acid, the possibility of its association with Reye's syndrome cannot be excluded.

Information for Patients

Patients should be informed of the following information before initiating therapy with an NSAID and periodically during the course of ongoing therapy. Patients should also be encouraged to read the NSAID Medication Guide that accompanies each prescription dispensed.

1. DOLOBID, like other NSAIDs, may cause serious CV side effects, such as MI or stroke, which may result in hospitalization and even death. Although serious CV events can occur without warning symptoms, patients should be alert for the signs and symptoms of chest pain, shortness of breath, weakness, slurring of speech, and should ask for medical advice when observing any indicative sign or symptoms. Patients should be apprised of the importance of this follow-up (see **WARNINGS, CARDIOVASCULAR EFFECTS**).
2. DOLOBID, like other NSAIDs, can cause GI discomfort and, rarely, serious GI side effects, such as ulcers and bleeding, which may result in hospitalization and even death. Although serious GI tract ulcerations and bleeding can occur without warning symptoms, patients should be alert for the signs and symptoms of ulcerations and bleeding, and should ask for medical advice when observing any indicative sign or symptoms including epigastric pain, dyspepsia, melena, and

hematemesis. Patients should be apprised of the importance of this follow-up (see **WARNINGS, Gastrointestinal Effects: Risk of Ulceration, Bleeding, and Perforation**).

3. DOLOBID, like other NSAIDs, can cause serious skin side effects such as exfoliative dermatitis, SJS, and TEN, which may result in hospitalizations and even death. Although serious skin reactions may occur without warning, patients should be alert for the signs and symptoms of skin rash and blisters, fever, or other signs of hypersensitivity such as itching, and should ask for medical advice when observing any indicative signs or symptoms. Patients should be advised to stop the drug immediately if they develop any type of rash and contact their physicians as soon as possible.
4. Patients should promptly report signs or symptoms of unexplained weight gain or edema to their physicians.
5. Patients should be informed of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, pruritus, jaundice, right upper quadrant tenderness, and “flu-like” symptoms). If these occur, patients should be instructed to stop therapy and seek immediate medical therapy.
6. Patients should be informed of the signs of an anaphylactic/anaphylactoid reaction (e.g. difficulty breathing, swelling of the face or throat). If these occur, patients should be instructed to seek immediate emergency help (see **WARNINGS**).
7. In late pregnancy, as with other NSAIDs, DOLOBID should be avoided because it may cause premature closure of the ductus arteriosus.

Laboratory Tests

Because serious GI tract ulcerations and bleeding can occur without warning symptoms, physicians should monitor for signs or symptoms of GI bleeding. Patients on long-term treatment with NSAIDs should have their CBC and a chemistry profile checked periodically. If clinical signs and symptoms consistent with liver or renal disease develop, systemic manifestations occur (e.g., eosinophilia, rash, etc.) or if abnormal liver tests persist or worsen, DOLOBID should be discontinued.

Drug Interactions

ACE-Inhibitors and Angiotensin II Antagonists

Reports suggest that NSAIDs may diminish the antihypertensive effect of ACE-inhibitors and angiotensin II antagonists. These interactions should be given consideration in patients taking NSAIDs concomitantly with ACE-inhibitors or angiotensin II antagonists. In some patients with compromised renal function, the co-administration of an NSAID and an ACE-inhibitor or an angiotensin II antagonist may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible.

Acetaminophen

In normal volunteers, concomitant administration of DOLOBID and acetaminophen resulted in an approximate 50% increase in plasma levels of acetaminophen. Acetaminophen had no effect on plasma levels of DOLOBID. Since acetaminophen in high doses has been associated with hepatotoxicity, concomitant administration of DOLOBID and acetaminophen should be used cautiously, with careful monitoring of patients.

Concomitant administration of DOLOBID and acetaminophen in dogs, but not in rats, at approximately 2 times the recommended maximum human therapeutic dose of each (40-52 mg/kg/day of DOLOBID/acetaminophen), resulted in greater gastrointestinal toxicity than when either drug was administered alone. The clinical significance of these findings has not been established.

Antacids

Concomitant administration of antacids may reduce plasma levels of DOLOBID. This effect is small with occasional doses of antacids, but may be clinically significant when antacids are used on a continuous schedule.

Aspirin

When DOLOBID is administered with aspirin, its protein binding is reduced, although the clearance of free DOLOBID is not altered. The clinical significance of this interaction is not known; however, as with other NSAIDs, concomitant administration of diflunisal and aspirin is not generally recommended because of the potential of increased adverse effects.

In normal volunteers, a small decrease in diflunisal levels was observed when multiple doses of DOLOBID and aspirin were administered concomitantly.

Cyclosporine

Administration of non-steroidal anti-inflammatory drugs concomitantly with cyclosporine has been associated with an increase in cyclosporine-induced toxicity, possibly due to decreased synthesis of renal prostacyclin. NSAIDs should be used with caution in patients taking cyclosporine, and renal function should be carefully monitored.

Diuretics

Clinical studies, as well as post marketing observations, have shown that DOLOBID can reduce the natriuretic effect of furosemide and thiazides in some patients. This response has been attributed to inhibition of renal prostaglandin synthesis.

In normal volunteers, concomitant administration of DOLOBID and hydrochlorothiazide resulted in significantly increased plasma levels of hydrochlorothiazide. DOLOBID decreased the hyperuricemic effect of hydrochlorothiazide. During concomitant therapy with NSAIDs, the patient should be observed closely for signs of renal failure (see **WARNINGS, Renal Effects**), as well as to assure diuretic efficacy.

Lithium

NSAIDs have produced an elevation of plasma lithium levels and a reduction in renal lithium clearance. The mean minimum lithium concentration increased 15% and the renal clearance was decreased by approximately 20%. These effects have been attributed to inhibition of renal prostaglandin synthesis by the NSAID. Thus, when NSAIDs and lithium are administered concurrently, subjects should be observed carefully for signs of lithium toxicity.

Methotrexate

NSAIDs have been reported to competitively inhibit methotrexate accumulation in rabbit kidney slices. This may indicate that they could enhance the toxicity of methotrexate. Caution should be used when NSAIDs are administered concomitantly with methotrexate.

NSAIDs

The administration of diflunisal to normal volunteers receiving indomethacin decreased the renal clearance and significantly increased the plasma levels of indomethacin. In some patients the combined use of indomethacin and DOLOBID has been associated with fatal gastrointestinal hemorrhage. Therefore, indomethacin and DOLOBID should not be used concomitantly.

The concomitant use of DOLOBID and other NSAIDs is not recommended due to the increased possibility of gastrointestinal toxicity, with little or no increase in efficacy. The following information was obtained from studies in normal volunteers.

Sulindac: The concomitant administration of DOLOBID and sulindac in normal volunteers resulted in lowering of the plasma levels of the active sulindac sulfide metabolite by approximately one-third.

Naproxen: The concomitant administration of DOLOBID and naproxen in normal volunteers had no effect on the plasma levels of naproxen, but significantly decreased the urinary excretion of naproxen and its glucuronide metabolite. Naproxen had no effect on plasma levels of DOLOBID.

Oral Anticoagulants

In some normal volunteers, the concomitant administration of DOLOBID and warfarin, acenocoumarol, or phenprocoumon resulted in prolongation of prothrombin time. This may occur because diflunisal competitively displaces coumarins from protein binding sites. Accordingly, when DOLOBID is administered with oral anticoagulants, the prothrombin time should be closely monitored during and for several days after concomitant drug administration. Adjustment of dosage of oral anticoagulants may be required. The effects of warfarin and NSAIDs on GI bleeding are synergistic, such that users of both drugs together have a risk of serious GI bleeding higher than users of either drug alone.

Tolbutamide

In diabetic patients receiving DOLOBID and tolbutamide, no significant effects were seen on tolbutamide plasma levels or fasting blood glucose.

Drug/Laboratory Test Interactions

Serum Salicylate Assays: Caution should be used in interpreting the results of serum salicylate assays when diflunisal is present. Salicylate levels have been found to be falsely elevated with some assay methods.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Diflunisal did not affect the type or incidence of neoplasia in a 105-week study in the rat given doses up to 40 mg/kg/day (equivalent to approximately 1.3 times the maximum recommended human dose), or in long-term carcinogenic studies in mice given diflunisal at doses up to 80 mg/kg/day (equivalent to approximately 2.7 times the maximum recommended human dose). It was concluded that there was no carcinogenic potential for DOLOBID.

Diflunisal passes the placental barrier to a minor degree in the rat. Diflunisal had no mutagenic activity after oral administration in the dominant lethal assay, in the Ames microbial mutagen test or in the V-79 Chinese hamster lung cell assay.

No evidence of impaired fertility was found in reproduction studies in rats at doses up to 50 mg/kg/day.

Pregnancy

Teratogenic Effects. Pregnancy Category C

A dose of 60 mg/kg/day of diflunisal (equivalent to two times the maximum human dose) was maternotoxic, embryotoxic, and teratogenic in rabbits. In three of six studies in rabbits, evidence of teratogenicity was observed at doses ranging from 40 to 50 mg/kg/day. Teratology studies in mice, at doses up to 45 mg/kg/day, and in rats at doses up to 100 mg/kg/day, revealed no harm to the fetus due to diflunisal. Aspirin and other salicylates have been shown to be teratogenic in a wide variety of species, including the rat and rabbit, at doses ranging from 50 to 400 mg/kg/day (approximately one to eight times the human dose). Animal reproduction studies are not always predictive of human response. There are no adequate and well controlled studies with diflunisal in pregnant women. DOLOBID should be used in pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nonteratogenic Effects

Because of the known effects of nonsteroidal anti-inflammatory drugs on the fetal cardiovascular system (closure of ductus arteriosus), use during pregnancy (particularly late pregnancy) should be avoided.

The known effects of drugs of this class on the human fetus during the third trimester of pregnancy include: constriction of the ductus arteriosus prenatally, tricuspid incompetence, and pulmonary hypertension; non-closure of the ductus arteriosus postnatally which may be resistant to medical management; myocardial degenerative changes, platelet dysfunction with resultant bleeding, intracranial bleeding, renal dysfunction or failure, renal injury/dysgenesis which may result in prolonged or permanent renal failure, oligohydramnios, gastrointestinal bleeding or perforation, and increased risk of necrotizing enterocolitis.

In rats at a dose of one and one-half times the maximum human dose, there was an increase in the average length of gestation. Similar increases in the length of gestation have been observed with aspirin, indomethacin, and phenylbutazone, and may be related to inhibition of prostaglandin synthetase.

Labor and Delivery

In rat studies with NSAIDs, as with other drugs known to inhibit prostaglandin synthesis, an increased incidence of dystocia, delayed parturition, and decreased pup survival occurred. The effects of DOLOBID on labor and delivery in pregnant women are unknown.

Nursing Mothers

Diflunisal is excreted in human milk in concentrations of 2-7% of those in plasma. Because of the potential for serious adverse reactions in nursing infants from DOLOBID, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness of DOLOBID in pediatric patients below the age of 12 have not been established. Use of DOLOBID in pediatric patients below the age of 12 is not recommended.

The adverse effects observed following diflunisal administration to neonatal animals appear to be species, age, and dose-dependent. At dose levels approximately 3 times the usual human therapeutic dose, both aspirin (200 to 400 mg/kg/day) and diflunisal (80 mg/kg/day) resulted in death, leukocytosis, weight loss, and bilateral cataracts in neonatal (4 to 5-day-old) beagle puppies after 2 to 10 doses. Administration of an 80 mg/kg/day dose of diflunisal to 25-day-old puppies resulted in lower mortality, and did not produce cataracts. In newborn rats, a 400 mg/kg/day dose of aspirin resulted in increased mortality and some cataracts, whereas the effects of diflunisal administration at doses up to 140 mg/kg/day were limited to a decrease in average body weight gain.

Geriatric Use

As with any NSAID, caution should be exercised in treating the elderly (65 years and older) since advancing age appears to increase the possibility of adverse reactions. Elderly patients seem to tolerate ulceration or bleeding less well than other individuals and many spontaneous reports of fatal GI events are in this population (see **WARNINGS, Gastrointestinal Effects – Risk of Ulceration, Bleeding, and Perforation**).

This drug is known to be substantially excreted by the kidney and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection and it may be useful to monitor renal function (see **WARNINGS, Renal Effects**).

ADVERSE REACTIONS

The adverse reactions observed in controlled clinical trials encompass observations in 2,427 patients.

Listed below are the adverse reactions reported in the 1,314 of these patients who received treatment in studies of two weeks or longer. Five hundred thirteen patients were treated for at least 24 weeks, 255 patients were treated for at least 48 weeks, and 46 patients were treated for 96 weeks. In general, the adverse reactions listed below were 2 to 14 times less frequent in the 1,113 patients who received short-term treatment for mild to moderate pain.

Incidence Greater Than 1%

Gastrointestinal

The most frequent types of adverse reactions occurring with DOLOBID are gastrointestinal: these include nausea**, vomiting, dyspepsia**, gastrointestinal pain**, diarrhea**, constipation, and flatulence.

Psychiatric

Somnolence, insomnia.

Central Nervous System

Dizziness.

Special Senses

Tinnitus.

Dermatologic

Rash**.

Miscellaneous

Headache**, fatigue/tiredness.

Incidence Less Than 1 in 100

The following adverse reactions, occurring less frequently than 1 in 100, were reported in clinical trials or since the drug was marketed. The probability exists of a causal relationship between DOLOBID and these adverse reactions.

Dermatologic

Erythema multiforme, exfoliative dermatitis, Stevens-Johnson syndrome, toxic epidermal necrolysis, urticaria, pruritus, sweating, dry mucous membranes, stomatitis, photosensitivity.

Gastrointestinal

Peptic ulcer, gastrointestinal bleeding, anorexia, eructation, gastrointestinal perforation, gastritis.

Liver function abnormalities; jaundice, sometimes with fever; cholestasis; hepatitis.

Hematologic

Thrombocytopenia; agranulocytosis; hemolytic anemia.

Genitourinary

Dysuria; renal impairment, including renal failure; interstitial nephritis; hematuria; proteinuria.

Psychiatric

Nervousness, depression, hallucinations, confusion, disorientation.

Central Nervous System

Vertigo; light-headedness; paresthesias.

Special Senses

Transient visual disturbances including blurred vision.

Hypersensitivity Reactions

Acute anaphylactic reaction with bronchospasm; angioedema; flushing.

Hypersensitivity vasculitis.

Hypersensitivity syndrome (see **WARNINGS, Hypersensitivity Syndrome**).

Miscellaneous

Asthenia, edema.

Causal Relationship Unknown

Other reactions have been reported in clinical trials or since the drug was marketed, but occurred under circumstances where a causal relationship could not be established. However, in these rarely reported events, that possibility cannot be excluded. Therefore, these observations are listed to serve as alerting information to physicians.

**Incidence between 3% and 9%. Those reactions occurring in 1% to 3% are not marked with an asterisk.

Respiratory

Dyspnea.

Cardiovascular

Palpitation, syncope.

Musculoskeletal

Muscle cramps.

Genitourinary

Nephrotic syndrome.

Special Senses

Hearing loss.

Miscellaneous

Chest pain.

A rare occurrence of fulminant necrotizing fasciitis, particularly in association with Group A β -hemolytic streptococcus, has been described in persons treated with non-steroidal anti-inflammatory agents, including diflunisal, sometimes with fatal outcome (see also **PRECAUTIONS, General**).

Potential Adverse Effects

In addition, a variety of adverse effects not observed with DOLOBID in clinical trials or in marketing experience, but reported with other non-steroidal analgesic/anti-inflammatory agents, should be considered potential adverse effects of DOLOBID.

OVERDOSAGE

Cases of overdosage have occurred and deaths have been reported. Most patients recovered without evidence of permanent sequelae. The most common signs and symptoms observed with overdosage were drowsiness, vomiting, nausea, diarrhea, hyperventilation, tachycardia, sweating, tinnitus, disorientation, stupor and coma. Diminished urine output and cardiorespiratory arrest have also been reported. The lowest dosage of DOLOBID at which a death has been reported was 15 grams without the presence of other drugs. In a mixed drug overdose, ingestion of 7.5 grams of DOLOBID resulted in death.

In the event of overdosage, the stomach should be emptied by inducing vomiting or by gastric lavage, and the patient carefully observed and given symptomatic and supportive treatment. Because of the high degree of protein binding, hemodialysis may not be effective.

The oral LD₅₀ of the drug is 500 mg/kg and 826 mg/kg in female mice and female rats respectively.

DOSAGE AND ADMINISTRATION

Carefully consider the potential benefits and risks of DOLOBID and other treatment options before deciding to use DOLOBID. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

After observing the response to initial therapy with DOLOBID, the dose and frequency should be adjusted to suit an individual patient's needs.

Concentration-dependent pharmacokinetics prevail when DOLOBID is administered; a doubling of dosage produces a greater than doubling of drug accumulation. The effect becomes more apparent with repetitive doses.

For mild to moderate pain, an initial dose of 1000 mg followed by 500 mg every 12 hours is recommended for most patients. Following the initial dose, some patients may require 500 mg every 8 hours.

A lower dosage may be appropriate depending on such factors as pain severity, patient response, weight, or advanced age; for example, 500 mg initially, followed by 250 mg every 8-12 hours.

For osteoarthritis and rheumatoid arthritis, the suggested dosage range is 500 mg to 1000 mg daily in two divided doses. The dosage of DOLOBID may be increased or decreased according to patient response.

Maintenance doses higher than 1500 mg a day are not recommended.

Tablets should be swallowed whole, not crushed or chewed.

HOW SUPPLIED

Tablets DOLOBID are capsule-shaped, film-coated tablets supplied as follows:

No. 3390 — 250 mg peach colored, coded DOLOBID on one side and MSD 675 on the other.

NDC 0006-0675-61 unit of use bottles of 60

(6505-01-164-0501, 250 mg 60's).

No. 3392 — 500 mg orange colored, coded DOLOBID on one side and MSD 697 on the other.

NDC 0006-0697-61 unit of use bottles of 60

(6505-01-144-9724, 500 mg 60's).

Distributed by:



MERCK & CO., INC., Whitehouse Station, NJ 08889, USA

Issued January 2007

Printed in USA

Medication Guide
for
Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)
(See the end of this Medication Guide for a list of prescription NSAID medicines.)

What is the most important information I should know about medicines called Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)?

NSAID medicines may increase the chance of a heart attack or stroke that can lead to death. This chance increases:

- with longer use of NSAID medicines
- in people who have heart disease

NSAID medicines should never be used right before or after a heart surgery called a “coronary artery bypass graft (CABG).”

NSAID medicines can cause ulcers and bleeding in the stomach and intestines at any time during treatment.

Ulcers and bleeding:

- can happen without warning symptoms
- may cause death

The chance of a person getting an ulcer or bleeding increases with:

- taking medicines called “corticosteroids” and “anticoagulants”
- longer use
- smoking
- drinking alcohol
- older age
- having poor health

NSAID medicines should only be used:

- exactly as prescribed
 - at the lowest dose possible for your treatment
 - for the shortest time needed
-

What are Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)?

NSAID medicines are used to treat pain and redness, swelling, and heat (inflammation) from medical conditions such as:

- different types of arthritis
- menstrual cramps and other types of short-term pain

Who should not take a Non-Steroidal Anti-Inflammatory Drug (NSAID)?

Do not take an NSAID medicine:

- if you had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAID medicine
- for pain right before or after heart bypass surgery

Tell your healthcare provider:

- about all of your medical conditions.
- about all of the medicines you take. NSAIDs and some other medicines can interact with each other and cause serious side effects. **Keep a list of your medicines to show to your healthcare provider and pharmacist.**

- if you are pregnant. **NSAID medicines should not be used by pregnant women late in their pregnancy.**
- if you are breastfeeding. **Talk to your doctor.**

What are the possible side effects of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)?

<p>Serious side effects include:</p> <ul style="list-style-type: none"> • heart attack • stroke • high blood pressure • heart failure from body swelling (fluid retention) • kidney problems including kidney failure • bleeding and ulcers in the stomach and intestine • low red blood cells (anemia) • life-threatening skin reactions • life-threatening allergic reactions • liver problems including liver failure • asthma attacks in people who have asthma 	<p>Other side effects include:</p> <ul style="list-style-type: none"> • stomach pain • constipation • diarrhea • gas • heartburn • nausea • vomiting • dizziness
---	---

Get emergency help right away if you have any of the following symptoms:

- shortness of breath or trouble breathing
- chest pain
- weakness in one part or side of your body
- slurred speech
- swelling of the face or throat

Stop your NSAID medicine and call your healthcare provider right away if you have any of the following symptoms:

- nausea
- more tired or weaker than usual
- itching
- your skin or eyes look yellow
- stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms and legs, hands and feet

These are not all the side effects with NSAID medicines. Talk to your healthcare provider or pharmacist for more information about NSAID medicines.

Other information about Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)

- Aspirin is an NSAID medicine but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines.
- Some of these NSAID medicines are sold in lower doses without a prescription (over-the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

NSAID medicines that need a prescription

Generic Name	Tradename
Celecoxib	Celebrex
Diclofenac	Cataflam, Voltaren, Arthrotec (combined with misoprostol)

Diflunisal	Dolobid
Etodolac	Lodine, Lodine XL
Fenoprofen	Nalfon, Nalfon 200
Flurbiprofen	Ansaid
Ibuprofen	Motrin, Tab-Profen, Vicoprofen* (combined with hydrocodone), Combunox (combined with oxycodone)
Indomethacin	Indocin, Indocin SR, Indo-Lemmon, Indomethegan
Ketoprofen	Oruvail
Ketorolac	Toradol
Mefenamic Acid	Ponstel
Meloxicam	Mobic
Nabumetone	Relafen
Naproxen	Naprosyn, Anaprox, Anaprox DS, EC-Naprosyn, Naprelan, Naprapac (copackaged with lansoprazole)
Oxaprozin	Daypro
Piroxicam	Feldene
Sulindac	Clinoril
Tolmetin	Tolectin, Tolectin DS, Tolectin 600

This Medication Guide has been approved by the U.S. Food and Drug Administration.

* Vicoprofen contains the same dose of ibuprofen as over-the-counter (OTC) NSAIDs, and is usually used for less than 10 days to treat pain. The OTC NSAID label warns that long term continuous use may increase the risk of heart attack or stroke.